

## **Product Datasheet**

Name: Mouse Anti-SARS-CoV-2 N protein Monoclonal Antibody

**Description:** Hybridoma clones have been derived from hybridization of myeloma cells with

spleen cells of BALB/c mouse immunized with recombinant SARS-CoV-2 N protein.

Catalog No.	Isotype	Clone No.	Usage	Buffer
bsm-41413M	lgG1	6A8	Capture /Detection	10mM PBS (pH7.4)

**Specificity:** Mab react with recombinant antigen SARS-CoV-2 N protein

**Host:** Mouse

Clonality: Monoclonal

Format: Liquid

**Concentration:** ≥1 mg/ml

**Purification:** ≥90% (SDS-PAGE)

**Preservative:** 0.1% Proclin300

**Application:** Recommended for sandwich immunoassays in ELISA and CLIA. Each laboratory

should determine an optimum working titer for use in its particular application.

**Storage:** Store at -20 °C for three years. Avoid repeated freeze/thaw cycles.

**Background:** Nucleocapsid protein is a most abundant protein of coronavirus. During virion assembly, N protein binds to viral RNA and leads to formation of the helical nucleocapsid. Nucleocapsid protein is a highly immunogenic phosphoprotein also implicated in viral genome replication and in modulating cell signaling pathways. Because of the conservation of N protein sequence and its strong immunogenicity, the N protein of coronavirus is chosen as a diagnostic tool.

**Note:** This product as supplied is intended for research or further manufacturing use only.




**400-901-9800** 

≥ sales@bioss.com.cn

support@bioss.com.cn

## **VALIDATION IMAGES**

## kDa M A B C D E F



Anti-SARS-CoV-2 N protein Mouse Monoclonal Antibody (Cat# bsm-41413M) at 1:1000 dilution.

Sample: rSARS-CoV-2 N Protein (Cat# bs-41408P)

Lane A: 50ng

Lane B: 25ng

Lane C: 5ng

Lane D: 2ng

Lane E: 1ng

Lane F: 0.5ng

Secondary

Goat Anti-Mouse IgG (H+L)/IRDye800CW at 1/20000 dilution.

Developed using the Odyssey (Li-cor).

Performed under reducing conditions.